

NOTE: Where map units are composed of multiple surficial materials, a compound map unit designator is used, separating more extensive materials from less extensive (e.g., Tb.Tv, Tb is more extensive than Tv).

QUATERNARY
HOLOCENE
NONGLACIAL ENVIRONMENT
ORGANIC DEPOSITS: Mostly saturated organic materials, consisting mainly of mosses, sedges, or other hydrophytic vegetation.
O Undifferentiated organic deposits: Bogs, fens, and swamps; generally occur where shallow lakes are infilled and in depressions along floodplains and abandoned meltwater channels.
COLLUVIAL DEPOSITS: Materials deposited by direct gravity-induced movement; lithologic composition dependent on source material, typically poorly sorted, massive to crudely stratified diamiction.
Cv Colluvial veneer: Deposits less <2 m thick that conform to underlying topography; typically on steep slopes.
LATE WISCONSINIAN
PROGLACIAL AND GLACIAL ENVIRONMENT
GLACIOACUSTRINE DEPOSITS: Well sorted, stratified sand, silt, and clay deposited by suspension settling and interflow and underflow currents; diamictions released from floating ice or colluviated from valley sides into glacial lakes; grain size may increase, and sorting may decrease in ice proximal environments.
GLv Glaciacustrine veneer: Deposits <2 m thick that conform to underlying topography; predominantly fine-grained material or silt-rich diamiction.
GLb Glaciacustrine blanket: Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
GLACIOFLUVIAL DEPOSITS: Poorly sorted to well sorted sand and gravel transported and deposited directly by glacial meltwater.
GFv Glaciofluvial veneer: Deposits <2 m thick that conform to underlying topography.
GFb Glaciofluvial blanket: Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography.
GFc Ice-contact deposits: Stratified sand and gravel with minor diamiction, forms hillocks and hollows.

TILL DEPOSITS: Diamictions eroded, transported and deposited at the base of an active glacier. They are dense, massive, and matrix supported and can be fissile and jointed. Matrix is typically composed of silt, sand, and clay. Clasts are often sub-angular to sub-rounded and can be faceted and striated.

Tv Till veneer: Deposits <2 m thick that conform to underlying topography; predominantly in upland regions with isolated bedrock exposures.

Tb Till blanket: Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.


Ts Streamlined till: Fillings, drumlins, and the sediment (down-ice) part of crag-and-tails.

ABLATION TILL DEPOSITS: Diamictions deposited by melt out, commonly from stagnant ice, of far-travelled supraglacial and englacial material. These deposits typically lack the density of basal till and have a high percentage of matrix sand. May be stratified and include sorted sands and gravels.

Tu Undulating till: Loose, sandy diamiction commonly representing thinner deposits near the margins of widespread ice stagnation, or in depressions where localized ice stagnation occurred; consist of hillocks and hollows with slopes ≤ 15°.

Th Hummocky till: Loose sandy diamiction commonly representing thicker deposits and widespread ice stagnation; consist of hillocks and hollows with slopes ≥ 15°.

PRE-QUATERNARY
BEDROCK: Lithology varies greatly across the map area and includes sedimentary, metamorphic, volcanic, and intrusive rocks of Permian to Quaternary age; outcrop is generally limited to areas of high relief.
R Undifferentiated bedrock: High-angle slopes in upland areas or in incised meltwater channels; may be susceptible to rock fall, hummocky, or undulating expressions are the result of glacial or meltwater erosion, or preferential erosion due to structural weaknesses; streamlined bedrock is the result of glacial erosion.



DESCRIPTIVE NOTES

The Satoh Mountain map area is in the Fraser Plateau, a physiographic subdivision of the Interior Plateau defined by a flat to gently rolling topography. Glacial sediments cover most of the region, and bedrock outcrops are rare (Holland, 1976). Previous work in the area includes glacial features mapping by Tipper (1971), To the east, Kerr and Giles (1993a, b) and Proudfoot and Allison (1993a, b) completed surficial geology mapping. Bedrock geology was originally mapped by Tipper (1969) and has been updated since by Bordet (2014) and Angen et al. (2017). The present basal till potential map continues the series published by Sacco et al. (2014a, b) for Geoscience BC's Targeting Resources for Exploration and Knowledge (TREK) project area (Clifford and Hart, 2014; Sacco et al., 2014a; Sacco and Jackaman, 2015).

Surficial sediment geochemical and mineralogical anomalies can be used to locate buried bedrock mineralization (Saarnisto, 1990; Klassen, 2001). Basal till is ideal for assessing bedrock-hosted mineral potential in areas covered by Quaternary sediments because it is commonly a first derivative of bedrock (Shilts, 1993), has a relatively simple and predictable transport history, and produces a geochemical and mineralogical signature that is more extensive than its bedrock source (Lewson, 2001). Glacial transport and deposition of basal till produces a dispersal train elongated down ice from its bedrock source (Fig. 1). To date, all till orientation surveys conducted in British Columbia have identified known mineralized sources (Plouffe et al., 2016).

The purpose of the basal till potential map series is to assist in the design of surface sediment exploration programs by identifying areas where basal till is most likely to occur. Ice flow indicators compiled by Arnold et al. (2016) are included in the maps to illustrate the general transport directions of basal till. These data should be supplemented with additional field measurements to assess local variability.

The basal till potential map series builds on earlier drift exploration potential maps developed by Proudfoot et al. (1995). Existing surficial geology, terrain, or soils and landform mapping data were reviewed and updated to produce the maps. Map unit definitions are based on conventions outlined by Cocking et al. (2016) and Deblonde et al. (2012) and unit colours are related to basal till potential classes. Each unit includes a label that describes the surficial material within it (mainly unconsolidated sediments) and its surface expression (individual plan-view forms and patterns of forms; Howes and Kenk, 1997).

New mapping focused on distinguishing basal till (Fig. 2) from ablation till (Fig. 3) which, because of a more complex transport and depositional history, is ill-suited for mineral exploration. The relationship between surface expression and till facies is predictable (Maynard, 1989; Aario and Peuraniemi, 1992; Proudfoot et al., 1995; Spirito et al., 2011; McClenaghan et al., 2013). For example, blanket, veneer, and streamlined units typically contain basal till facies, whereas undulating and hummocky units typically contain ablation till facies. Based on these relationships we used air photographs supplemented by sparse field data to construct our maps.

Basal till consists of sediment eroded, transported and deposited at the base of an active glacier (Dreimanns, 1988). It typically has a relatively subparallel surface expression that either follows underlying topography (Tb, Tv) or is streamlined in the direction of ice flow (Ts). It is a dense, massive, matrix supported diamicton, with a matrix mainly of silt with lesser

