


Basal till potential of the Carnlick Creek map area (NTS 093C/14), British Columbia

D. Sacco, H. Arnold, T. Ferbey, and W. Jackaman



- NOTE: Where map units are composed of multiple surficial materials, a compound map unit designator is used, separating more extensive materials from less extensive (e.g., for Tb, Tv, Th is more extensive than Tv).
- QUATERNARY**
- HOLOCENE**
- NONGLACIAL ENVIRONMENT**
- ORGANIC DEPOSITS:** Mostly saturated organic materials, consisting mainly of mosses, sedges, or other hydrophytic vegetation.
- O Undifferentiated organic deposits:** Bogs, fens, and swamps; generally occur where shallow lakes are infilled and in depressions along floodplains and abandoned meltwater channels.
 - Colluvial veneer:** Deposits less <2 m thick that conform to underlying topography; typically on steep slopes.
 - Colluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
- LATE WISCONSINAN**
- PROGLACIAL AND GLACIAL ENVIRONMENT**
- GLACIOFLUVIAL DEPOSITS:** Well sorted, stratified sand, silt, and clay deposited by suspension setting and interflow and underflow currents; diamictons released from floating ice or colluviated from valley sides into glacial lakes; grain size may increase, and sorting may decrease in ice proximal environments.
- GLv Glaciofluvial veneer:** Deposits <2 m thick that conform to underlying topography; predominantly fine-grained material or silt-rich diamictons.
 - GLb Glaciofluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
 - GLv Glaciofluvial veneer:** Deposits <2 m thick that conform to underlying topography.
 - GLb Glaciofluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography.
- GLACIOFLUVIAL DEPOSITS:** Poorly sorted to well sorted sand and gravel transported and deposited directly by glacial meltwater.
- Gfv Glaciofluvial veneer:** Deposits <2 m thick that conform to underlying topography.
 - Gfb Glaciofluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography.
- GFC Ice-contact deposits:** Stratified sand and gravel with minor diamicton deposited, forms hillocks and hollows.
- GFR Eskers:** Sinuous ridges of stratified sand and gravel deposited in subglacial, englacial, or supraglacial channels.
- BASAL TILL DEPOSITS:** Diamictons eroded, transported and deposited at the base of an active glacier. They are dense, massive, and matrix supported and can be fissile and jointed. Matrix is typically composed of silt, sand, and clay. Clasts are often sub-angular to sub-rounded and can be faceted and striated.
- Tv Till veneer:** Deposits <2 m thick that conform to underlying topography; predominantly in upland regions with isolated bedrock exposures.
 - Tb Till blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
 - Ts Streamlined till:** Flutings, drumlins, and the sediment (down-ice) part of crag-and-tails.
 - ABLATION TILL DEPOSITS:** Diamictons deposited by melt out, commonly from stagnation ice, of far-travelled supraglacial and englacial material. These deposits typically lack the density of basal till and have a high percentage of matrix sand. May be stratified and include sorted sands and gravels.
 - Tu Undulating till:** Loose, sandy diamicton commonly representing thinner deposits near the margins of widespread ice stagnation, or in depressions where localized ice stagnation occurred; consist of hillocks and hollows with slopes $\leq 15^\circ$.
 - Th Hummocky till:** Loose sandy diamicton commonly representing thicker deposits and widespread ice stagnation; consist of hillocks and hollows with slopes $\leq 15^\circ$.
- OTHER TILL DEPOSITS:** Diamictons composed of subglacial, englacial, and (or) supraglacial debris produced by glacial thrust, push, or meltout processes.
- Tr Ridged till:** Elongate ridges oriented perpendicular to ice-flow direction, formed subglacially or at glacial margin.
 - Undifferentiated bedrock:** High-angle slopes in upland areas or in incised meltwater channels; may be susceptible to rock fall; hummocky or undulating expressions are the result of glacial or meltwater erosion, or preferential erosion due to structural weaknesses; streamlined bedrock is the result of glacial erosion.

DESCRIPTIVE NOTES

The Carnlick Creek map area is in the Nechako and Fraser Plateaus, physiographic subdivisions of the Interior Plateau defined by a flat to gently rolling topography. Glacial sediments cover much of the region, and bedrock outcrops are rare (Holland, 1973). Previous work in the area includes soils and terrain mapping by Baender (1980) and glacial features mapping by Tipper (1971). To the east, Kerr and Giles (1993a, b) and Prud'homme and Allison (1993a, b) completed surficial geology mapping. Bedrock geology was originally mapped by Tipper (1967) and has been updated since by Southern and Southern (1984), Bordet (2014), and Angen et al. (2017). The present basal till potential map continues the series published by Sacco et al. (2014a to j) for Geoscience BC's Targeting Resources for Exploration and Knowledge (TREG) project area (Clifford and Hart, 2014; Sacco et al., 2014k; Sacco and Jackaman, 2015).

Surficial sediment geology and mineralogical anomalies can be used to locate buried bedrock mineral potential in areas covered by Quaternary sediments because it is commonly a first derivative of bedrock (Shilts, 1993), has a relatively simple and predictable transport history, and produces a geochemical and mineralogical signature that is more extensive than its bedrock source (Levon, 2001). Glacial transport and deposition of basal till produces a dispersal train elongated down ice from its bedrock source (Fig. 1). To date, all tilt orientation surveys conducted in British Columbia have identified known mineralized sources (Plouffe et al., 2016).

The purpose of the basal till potential map series is to assist in the design of surface sediment exploration programs by identifying areas where basal till is most likely to occur. Ice flow indicators compiled by Arnold et al. (2016) are included in the maps to illustrate the general transport directions of basal till. These data should be supplemented with additional field measurements to assess for local variability.

In a basal till potential map, each unit with till, as a primary or secondary surface material is assigned a basal till potential rating. High potential is assigned to units containing mainly basal till. The highest potential category (1) includes till blankets (>2 m thick) and streamlined till with some till veneer (<2 m thick). In these units, samples can be collected from most exposures. In the second category of high potential (2), till veneers are predominant and likely include some bedrock exposures. In these areas, sample collection may be most productive down-ice from bedrock outcrop, where till might be sufficiently thick to avoid post-depositional surface processes such as pedogenesis. In the third category of high potential (3), map units are mostly basal till (Tb, Ts, Tv) with lesser amounts of another surface material (excluding Tu or Th). Knowledge of the surface expression of this secondary material, which is provided in the map unit label, will assist in targeting basal till.

Moderate potential is assigned to units containing varying amounts of basal till and ablation till. These map units typically represent (4) thick basal till deposits in depressions or small valleys where ablation till has been deposited, or (5) near the margins of extensive ablation till map units where basal till may be found within a few metres of surface or in areas of higher elevation where ablation till thins.

Low potential (6) is assigned to units consisting mainly of surface material other than till. These areas may include basal till deposits that are too small to resolve at the current map scale, or are discontinuous. Poor potential (7) is assigned to areas of thick ablation till. These areas typically consist of hummocky ablation till and may include lesser amounts of another surficial material (e.g., ice-contact glaciofluvial deposits). These areas are still mapped as having potential because underlying basal till deposits may be present at depth.

Acknowledgements

We thank L.B. Aspler and A.S. Hickin for reviewing this map. The British Columbia Ministry of Forests, Lands and Natural Resource Operations provided access to stereo models and imagery.

References cited

Angen, J.J., Rattini, M., Hart, C.J.R., Westberg, E., Logan, J.M., and Kim, R., 2017. Bedrock geology, TREG project area, northern Interior Plateau, central British Columbia. Geoscience BC Map 2017-06-01 and WDRU Map 12-2017, 1:250,000 scale.

Arnold, H., Ferbey, T., and Hickin, A.S., 2016. Ice-flow indicator compilation, British Columbia and Yukon. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2016-04. Geological Survey of Canada Open File 8983, 1:1,500,000 scale.

Baender, H., 1980. Upper Dean River resource folio, terrain. British Columbia Ministry of Environment, unpublished, 1:50,000 scale.

Bordet, E., 2014. Eocene volcanic response to the tectonic evolution of the Canadian Cordillera. Unpublished Ph.D. thesis, University of British Columbia, 488 p.

Clifford, A., and Hart, C.J.R., 2014. Targeting Resources through Exploration and Knowledge (TREG): Geoscience BC's newest minerals project, Interior Plateau region, central British Columbia (NTS 093B, C, F, G). In: Geoscience BC Summary of Activities 2013, Geoscience BC, Report 2014-1, pp. 13-18.

Cocking, R.D., Deblonde, C., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Laviolette, A., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2016. Surficial geology model, version 2.2.0: Revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps. Geological Survey of Canada, Open File 8941, 45 p.

Deblonde, C., Plouffe, A., E. Boisvert-Buller, G., Davenport, P., Everett, D., Huntley, D., Inglis, E., Kerr, D., Moore, A., Paradis, S.J., Parent, M., Smith, I.R., St. Onge, D., and Weatherston, A., 2012. Science language for an integrated Geological Survey of Canada data model for surficial geology maps, version 1.2. Geological Survey of Canada, Open File 7003, 224 p.

Dreimanis, A., 1989. Tills: their genetic terminology and classification. In: Goldthwait, R.P., and Matsch, C.L. (Eds.), Genetic Classification of Glacial Deposits, Balkema, Rotterdam, 17-83.

Ferbey, T., and Levon, V.M., 2009. The influence of ice-flow reversals on the vertical and horizontal distribution of trace elements in tills. Huckleberry mine area, west-central British Columbia. In: Paulen, R.C., and McMartin, I. (Eds.), Application of Till and Stream Sediment Heavy Mineral and Geochemical Methods to Mineral Exploration in Western and Northern Canada, Geological Association of Canada, Short Course Notes 18, pp. 145-151.

Holland, S.B., 1976. Landforms of British Columbia: a physiographic outline. British Columbia Ministry of Energy and Mines, and Petroleum Resources, British Columbia Geological Survey Bulletin 46, 138 p.

Howes, D.E., and Kenk, E., 1997. Terrain classification system for British Columbia (revised Edition). British Columbia Ministry of Environment, Manual 10 (Version 2), 102 p.

Jackaman, W., Sacco, D.A., and Lett, R.E., 2015. Regional geochemical and mineralogical data, TREG project - year 2 Interior Plateau, British Columbia. Geoscience BC, Report 2015-12, 13 p.

Kerr, D.E., and Giles, T.R., 1993a. Surficial geology of the Chilako Forks area (NTS 93C/1). British Columbia Ministry of Energy and Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-16, 1:50,000 scale.

Kerr, D.E., and Giles, T.R., 1993b. Surficial geology of the Cheekat area (NTS 93C/8). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-17, 1:50,000 scale.

Klassen, R.A., 2001. A Quaternary geological perspective on geochemical exploration in glaciated terrain. In: McClenghan, M.B., Bobrowsky, P.T., Hall, G.E.M., and Cook, S.J. (Eds.), Drift Exploration in Glaciated Terrain, Geological Society, Special Publication 185, p. 1-17.

Lefebvre, D.V., and Ray, G.E. (Eds.), 1995. Selected British Columbia mineral deposit profiles, volume 1 - metallic and coal. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1995-20, digital.

Lefebvre, D.V., and Hoot, T., 1990. Selected British Columbia mineral deposit profiles, volume 2 - more metallic deposits. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1990-13, digital.

Levon, V.M., 2001. Regional till geochemical surveys in the Canadian Cordillera: sample media, methods, and anomaly evaluation. In: McClenghan, M.B., Bobrowsky, P.T., Hall, G.E.M., and Cook, S.J. (Eds.), Drift Exploration in Glaciated Terrain, Geological Society, Special Publication 185, p. 45-68.

Maynard, D.E., 1989. Terrain analysis in exploration geochemistry. Exploire, 65, 14-16.

McClenghan, M.B., Plouffe, A., McMartin, I., Campbell, J.E., Spirito, V.A., Paulen, R.C., Garrett, R.G., Hall, G.E.M., 2013. Till sampling and geochemical analytical protocols used by the Geological Survey of Canada. Geoscience: Exploration, Environment, Analysis, 13, 285-301.

Miller, J.K., 1984. Model for clastic indicator trains in till. In: Prospecting in areas of glaciated terrain. Institution of Mining and Metallurgy, London, pp. 69-77.

Plouffe, A., Ferbey, T., Hashmi, S., and Ward, B.C., 2016. Till geochemistry and mineralogy: vectoring towards Cu porphyry deposits in British Columbia, Canada. Geochemistry: Exploration, Environment, Analysis, <http://doi.org/10.1144/gechem-2015-398>.

Prud'homme, D.N., and Allison, R.F., 1993a. Surficial geology of the Clusko River area (NTS 93C/9). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-18, 1:50,000 scale.

Prud'homme, D.N., and Allison, R.F., 1993b. Surficial geology of the Toll Mountain area (NTS 93C/16). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-19, 1:50,000 scale.

Prud'homme, D.N., Bobrowsky, P.T., and Meldrum, G.O., 1985. Drift exploration potential maps derived from terrain geology maps. In: Bobrowsky, P.T., Sibick, S.L., Newell, J.M., and Mayska, P.F. (Eds.), Drift Exploration in the Canadian Cordillera, British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Paper 1985-2, p. 23-31.

Sacco, D.A., and Jackaman, W., 2015. Targeted geochemical and mineralogical surveys in the TREG project area, central British Columbia (parts of NTS 093B, C, F, G), year two. In: Geoscience BC Summary of Activities 2014, Geoscience BC, Report 2015-1, pp. 1-12.

Sacco, D., Ferbey, T., and Jackaman, W., 2014a. Basal till potential of the Clusko River map area (NTS 093C/9). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-06. Geoscience BC Map 2014-06-01, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014b. Basal till potential of the Mamont Lake map area (NTS 093B/12). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-07. Geoscience BC Map 2014-06-02, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014c. Basal till potential of the Marmot Lake map area (NTS 093B/13). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-08. Geoscience BC Map 2014-06-03, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014d. Basal till potential of the Toll Mountain map area (NTS 093C/16). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-09. Geoscience BC Map 2014-06-04, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014e. Basal till potential of the Suscha Creek map area (NTS 093A/1). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-10. Geoscience BC Map 2014-06-05, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014f. Basal till potential of the Cogitako River map area (NTS 093G/04). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-11. Geoscience BC Map 2014-06-06, 1:50,000 scale.

Basal till potential

High

- 1 Only thick basal till (Tb or Ts); may contain lesser amounts of thin basal till (e.g., Tb, Tv).
- 2 Only thin basal till (Tv); may contain lesser amounts of thick basal till (e.g., Tv, Tb).
- 3 Basal till with lesser amounts of another surficial material, excluding ablation till (e.g., Tb, Cb, Tv, R).

Moderate

- 4 Basal till with lesser amounts of ablation till (e.g., Tb, Tu).
- 5 Ablation till with lesser amounts of basal till (e.g., Tu, Tb).

Low

- 6 Another surficial material with lesser amounts of basal till (e.g., Cb, Tb).

Poor

- 7 Only ablation or ridged till at surface (Th, Tu, or Tr) or ablation or ridged till and another surficial material, excluding basal till (e.g., Th, O; Gfb, Tu; Tu, Tr).

None

- Surficial material other than till.

TILL SAMPLES (Labeled with sample number; Jackaman et al., 2015)

Matrix geochemistry
Geochemistry and mineralogy.

ICE-FLOW INDICATORS (Arnold et al., 2016)

Crag-and-tail (flow direction known)
Drumlin, Drumlinoid or fluting (flow direction known, unknown)
Striation (flow direction known)

National and Provincial parks and protected areas
Road

MINERAL OCCURRENCES

MINFILE NUMBER NAME STATUS COMMODITY DEPOSIT TYPE*

No MINFILE occurrences in map area, as of November 2017.

*If deposit type is available, see Lefebvre and Ray (1995), Lefebvre and Hoy (1996), and Simandl et al. (1999) for mineral deposit profile codes and definitions.

PLAN VIEW

CROSS SECTIONS

LONGITUDINAL SECTION

Fig. 1. Model of clastic dispersal in basal till (modified from Miller, 1984). Highest values (dark blue) define the head of a dispersal train at the head, and decrease exponentially in the direction of ice flow (Tv). It is a dense, massive, matrix-supported till.

Fig. 2. Basal till in vertical exposure. Note blocky appearance. Granule and coarser-sized clasts float in a clay-silt matrix. Measuring tape in centimetres.

Fig. 3. Ablation till exposed in road cut with higher percentage of sand and gravel, and lower density, compared to typical basal till (see Fig. 2). Pick with scale (65 cm).

Fig. 4. Map showing the location of the Carnlick Creek map area (NTS 093C/14) in British Columbia, with an inset map of the region.

Table of Mineral Occurrences:

MINFILE NUMBER	NAME	STATUS	COMMODITY	DEPOSIT TYPE*
093F04	QUALCHO LAKE			
093F03	FAMNE CREEK			
093G02	TEACH LAKE			
B052 of 2017-06				
GBC Map 2017-02-05				
093G01	KAROLFO			
093G14	CARLICK CREEK			
093G15	KUSHYA RIVER			
B055 of 2017-05				
GBC Map 2017-02-04				
093G16				
B056 of 2017-04				
GBC Map 2017-02-03				
093G18	CHRISTENSEN CREEK			
093G19	DOWNTOWN CREEK			
093G20				