


## Basal till potential of the Kushya River map area (NTS 093C/15), British Columbia

D. Sacco, H. Arnold, T. Ferbey, and W. Jackaman



**NOTE:** Where map units are composed of multiple surficial materials, a compound map unit designator is used, separating more extensive materials from less extensive (e.g., for Tb.Tv, Tb is more extensive than Tv).

- QUATERNARY**
- HOLOGENE**
- NONGLACIAL ENVIRONMENT**
- ORGANIC DEPOSITS:** Mostly saturated organic materials, consisting mainly of mosses, sedges, or other hydrophytic vegetation.
- O Undifferentiated organic deposits:** Bogs, fens, and swamps; generally occur where shallow lakes are infilled and in depressions along floodplains and abandoned meltwater channels.
- COLLUVIAL DEPOSITS:** Materials deposited by direct gravity-induced movement; lithologic composition dependent on source material; typically poorly sorted, massive to crudely stratified diamict.
- Cv Colluvial veneer:** Deposits less <2 m thick that conform to underlying topography, typically on steep slopes.
  - Cb Colluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; typically on steep slopes.
- LATE WISCONSINAN**
- PROGLACIAL AND GLACIAL ENVIRONMENT**
- GLACIOFLUVIAL DEPOSITS:** Poorly sorted to well sorted sand and gravel transported and deposited directly by glacial meltwater.
- Gfv Glaciofluvial veneer:** Deposits <2 m thick that conform to underlying topography.
  - Gfb Glaciofluvial blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography.
  - Gfc Ice-contact deposits:** Stratified sand and gravel with minor diamict, forms hillocks and hollows.
  - Gfr Eskers:** Sinuous ridges of stratified sand and gravel deposited in subglacial, englacial, or supraglacial channels.

- BASAL TILL DEPOSITS:** Diamictions eroded, transported and deposited at the base of an active glacier. They are dense, massive, and matrix supported and can be fissile and jointed. Matrix is typically composed of silt, sand, and clay. Clasts are often sub-angular to sub-rounded and can be faceted and striated.
- Tv Till veneer:** Deposits <2 m thick that conform to underlying topography; predominantly in upland regions with isolated bedrock exposures.
  - Tb Till blanket:** Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
  - Ts Streamlined till:** Flutings, drumlins, and the sediment (down-ice) part of crag-and-tails.
- ABLATION TILL DEPOSITS:** Diamictions deposited by melt out, commonly from stagnant ice, of far-travelled supraglacial and englacial material. These deposits typically lack the density of basal till and have a high percentage of matrix sand. May be stratified and include sorted sands and gravels.
- Tu Undulating till:** Loose, sandy diamict commonly representing thinner deposits near the margins of widespread ice stagnation, or in depressions where localized ice stagnation occurred; consist of hillocks and hollows with slopes  $\leq 15^\circ$ .
  - Th Hummocky till:** Loose sandy diamict commonly representing thicker deposits and widespread ice stagnation; consist of hillocks and hollows with slopes  $\leq 15^\circ$ .

### Basal till potential

- High**
  - 1 Only thick basal till (Tb or Ts); may contain lesser amounts of thin basal till (e.g., Tb.Tv).
  - 2 Only thin basal till (Tv); may contain lesser amounts of thick basal till (e.g., Tv.Tb).
  - 3 Basal till with lesser amounts of another surficial material, excluding ablation till (e.g., Tb.Cb, Tv.R).
- Moderate**
  - 4 Basal till with lesser amounts of ablation till (e.g., Tb.Tu).
  - 5 Ablation till with lesser amounts of basal till (e.g., Tu.Tb).
- Low**
  - 6 Another surficial material with lesser amounts of basal till (e.g., Cb.Tb).
- Poor**
  - 7 Only ablation or ridged till at surface (Th.Tu, or Tr), or ablation or ridged till and another surficial material, excluding basal till (e.g., Th.O, Gfb.Tv, Tu.Tr).
- None**
  - Surficial material other than till.

**TILL SAMPLES (Labeled with sample number; Jackaman et al., 1915)**

Matrix geochemistry and mineralogy.

**ICE-FLOW INDICATORS (Arnold et al., 2016)**

Crag-and-tail (flow direction known)  
Drumlin, Drumlinoid or fluting (flow direction known, unknown).

National and Provincial parks and protected areas  
Road

MINFILE NUMBER	NAME	STATUS	COMMODITY	DEPOSIT TYPE*
----------------	------	--------	-----------	---------------

No MINFILE occurrences in map area, as of November 2017.  
\*If deposit type is available, see LeBlond and Ray (1995), LeBlond and Hoy (1996), and Simard et al. (1999) for mineral deposit profile codes and definitions.

### DESCRIPTIVE NOTES

The Kushya River map area is in the Nechako and Fraser Plateaus, physiographic subdivisions of the Interior Plateau defined by a flat to gently rolling topography. Glacial sediments cover much of the region, and bedrock outcrops are rare (Holland, 1976). Previous work in the area includes glacial features mapping by Tipper (1971), to the east, Kerr and Giles (1993a, b) and Proudfoot and Allison (1993a, b) completed surficial geology mapping. Bedrock geology was originally mapped by Tipper (1969) and has been updated since by Bordet (2014) and Argen et al. (2017). The present basal till potential map continues the series published by Sacco et al. (2014a to j) for Geoscience BC's Targeting Resources for Exploration and Knowledge (TREK) project area (Clifford and Hart, 2014; Sacco et al., 2014k; Sacco and Jackaman, 2015).

Surficial sediment geochemical and mineralogical anomalies can be used to locate buried bedrock mineralization (Saarnisto, 1990; Klassen, 2001). Basal till is ideal for assessing bedrock-hosted mineral potential in areas covered by Quaternary sediments because it is commonly a first derivative of bedrock (Shilts, 1993), has a relatively simple and predictable transport history, and produces a geochemical and mineralogical signature that is more extensive than its bedrock source (Leveson, 2001). Glacial transport and deposition of basal till produces a dispersal train elongated down ice from its bedrock source (Fig. 1). To date, all till orientation surveys conducted in British Columbia have identified known mineralized sources (Proudfoot et al., 2016).

The purpose of the basal till potential map series is to assist in the design of surface sediment exploration programs by identifying areas where basal till is most likely to occur. Ice flow indicators compiled by Arnold et al. (2016) are included in the map to illustrate the general transport directions of basal till. These data should be supplemented with additional field measurements to assess for local variability.

The basal till potential map series builds on earlier drift exploration potential maps developed by Proudfoot et al. (1995). Existing surficial geology, terrain, or soils and landform mapping data were reviewed and updated to produce the maps. Map unit definitions are based on conventions outlined by Cocking et al. (2016) and Deblonde et al. (2012) and unit colours are related to basal till potential classes. Each unit includes a label that describes the surficial material within it (mainly basal till (sediments) and its surface expression (individual plan-view forms and patterns of forms; Howes and Kenk, 1997).

New mapping focused on distinguishing basal till (Fig. 2) from ablation till (Fig. 3) which, because of a more complex transport and depositional history, is ill-suited for mineral exploration. The relationship between surface expression and till facies is predictable (Maynard, 1989; Aario and Peuranieni et al., 2013). For example, blanket, veneer, and streamlined units typically contain basal till facies, whereas undulating and hummocky units typically contain ablation till facies. Based on these relationships we used air photographs supplemented by sparse field data to construct our maps.

Basal till consists of sediment eroded, transported, and deposited at the base of an active glacier (Dreimann, 1989). It typically has a relatively subrounded surface expression that either follows underlying topography (Tb, Tv) or is streamlined in the direction of ice flow (Ts). It is dense, massive, matrix-supported diamict, with a matrix mainly of silt with lesser

amounts of sand and clay (Fig. 2). Vertical joints and subhorizontal fissility intersections can give basal till a blocky appearance in section. Clasts are mostly subangular to subrounded and are commonly striated. The transport path of basal till is relatively simple and short and can be established by measuring the azimuth of ice-flow indicators produced by subglacial flow. However, multiple ice-flow events can create a more complex transport path, highlighting the importance of ice-flow history reconstructions (Ferbey and Leveson, 2009; Proudfoot et al., 2016).


Compared to basal till, the transport distance of ablation till is longer and the depositional history more complex. Ablation till consists of material transported in the englacial and supraglacial environments and commonly deposited by passive melt out processes. Melting of remnant ice-blocks mantled or surrounded by glacial debris produces irregular, undulating to hummocky topography (Tu, Th). Ablation till is less consolidated, has a higher percentage of gravel-sized material and a sandier matrix (Fig. 3). It can be massive to crudely stratified and may contain lenses of sorted sand and gravel. Deposited during deglaciation, ablation tills are typically the youngest Late Wisconsinan till facies exposed at surface, and can overlie basal tills. Windows through an ablation till, into an underlying basal till, can exist but may be indistinguishable in air photographs.

In a basal till potential map, each unit with till, as a primary or secondary surface material is assigned a basal till potential rating. High potential is assigned to units containing mainly basal till. The highest potential category (1) includes till blankets (>2 m thick) and streamlined till with some till veneer (<2 m thick). In these units, samples can be collected from most exposures. In the second category of high potential (2), till veneers are predominant and likely include some bedrock exposures. In these areas, sample collection may be most productive down-ice from bedrock outcrop, where till might be sufficiently thick to avoid post-depositional surface processes such as pedogenesis. In the third category of high potential (3), map units are mostly basal till (Tb, Tv) with lesser amounts of another surface material (excluding Tu or Th). Knowledge of the surface expression of this secondary material, which is provided in the map unit label, will assist in targeting basal till.

Moderate potential is assigned to units containing varying amounts of basal till and ablation till. These map units typically represent (4) thick basal till deposits in depressions or small valleys where ablation till has been deposited, or (5) near the margins of extensive ablation till map units where basal till may be found within a few metres of surface or in areas of higher elevation where ablation till thins.

Low potential (6) is assigned to units consisting mainly of surface material other than till. These areas may include basal till deposits that are too small to resolve at the current map scale, or are discontinuous. Poor potential (7) is assigned to areas of thick ablation till. These areas typically consist of hummocky ablation till and may include lesser amounts of another surficial material (e.g., ice-contact glaciofluvial deposits). These areas are still mapped as having potential because underlying basal till deposits may be present at depth.

**Acknowledgements**  
We thank L.B. Aspler and A.S. Harkin for reviewing this map. The British Columbia Ministry of Energy, Mines and Natural Resource Operations provided access to stereo models and imagery.



References cited

Aario, R., and Peuranieni, V., 1992. Glacial dispersal of till constituents in moraine landforms of different types. *Geomorphology* 6, 9-25.

Argen, J.J., Rattim, M., Hart, C.J.R., Westberg, E., Logan, J.M., and Kim, R., 2017. Bedrock geology, TREK project area, northern Interior Plateau, central British Columbia. Geoscience BC Map 2017-06-01 and MDU Map 12-2017, 1:250,000 scale.

Arnold, H., Ferbey, T., and Hickin, A.S., 2016. Ice-flow indicator compilation, British Columbia and Yukon. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2016-04, Geological Survey of Canada Open File 9083, 1:1,750,000 scale.

Bordet, E., 2014. Eocene volcanic response to the tectonic evolution of the Canadian Cordillera. Unpublished Ph.D. thesis, University of British Columbia, 486 p.

Clifford, A., and Hart, C.J.R., 2014. Targeting Resources through Exploration and Knowledge (TREK): Geoscience BC's newest minerals project, Interior Plateau region, central British Columbia (NTS 093C). C. F. G. In: Geoscience BC Summary of Activities 2013, Geoscience BC, Report 2014-1, pp. 13-16.

Cocking, R.B., Deblonde, C., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Laviolette, A., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2016. Surficial data model, version 2.2.0. Revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps. Geological Survey of Canada, Open File 9041, 45 p.

Deblonde, C., Plouffe, A., E. Boisvert, Butler, G., Davenport, P., Everett, D., Huntley, D., Inglis, E., Kerr, D.,

Moore, A., Paradis, S.J., Parent, M., Smith, I.R., St. Onge, D., and Weatherston, A., 2012. Science language for an integrated Geological Survey of Canada data model for surficial geology maps, version 1.2. Geological Survey of Canada, Open File 7003, 224 p.

Dreimann, A., 1989. Tills: their genetic terminology and classification. In: Goldswirth, R.P., and Matsch, C.L. (Eds.), *Genetic Classification of Glacio-genetic Deposits*, Balkema, Rotterdam, 17-83.

Ferbey, T., and Leveson, V.M., 2009. The influence of ice-flow reversal on the vertical and horizontal distribution of trace elements in tills, Huckleberry mine area, west-central British Columbia. In: Paulen, R.C., and McMartin, I. (Eds.), *Application of Till and Stream Sediment Heavy Mineral and Geochemical Methods to Mineral Exploration in Western and Northern Canada*, Geological Association of Canada, Short Course Notes 18, pp. 145-151.

Holland, S.S., 1976. Landforms of British Columbia: a physiographic outline. British Columbia Ministry of Energy and Mines, and Petroleum Resources, British Columbia Geological Survey Bulletin 48, 138 p.

Howes, D.E., and Kenk, E., 1997. Terrain classification system for British Columbia (revised Edition). British Columbia Ministry of Environment, Manual 10 (Version 2), 102 p.

Jackaman, W., Sacco, D.A., and Lett, R.E., 2015. Regional geochemical and mineralogical data, TREK project-year 2, Interior Plateau, British Columbia. Geoscience BC, Report 2015-12, 13 p.

Kerr, D.E., and Giles, T.R., 1993a. Surficial geology of the Chilikeo Forks area (NTS 82C/1). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-16, 1:50,000 scale.

Kerr, D.E., and Giles, T.R., 1993b. Surficial geology of the Chezacat area (NTS 93C/8). British Columbia

Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-17, 1:50,000 scale.

Klassen, R.A., 2001. A Quaternary geological perspective on geochemical exploration in glaciated terrain. In: McEneaney, M.B., Bobrowsky, P.T., Stribick, S.J., and Cook, S.J. (Eds.), *Drift Exploration in Glaciated Terrain*, Geological Society, Special Publication 195, p. 1-17.

LeBlond, D.V., and Hoy, T. (Eds.), 1996. Selected British Columbia mineral deposit profiles, volume 1 - metallics and coal. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1996-20, digital.

LeBlond, D.V., and Hoy, T. (Eds.), 1996. Selected British Columbia mineral deposit profiles, volume 2 - more metallic deposits. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1996-13, digital.

Leveson, V.M., 2001. Regional till geochemical surveys in the Canadian Cordillera: sample media, methods, and anomaly evaluation. In: McEneaney, M.B., Bobrowsky, P.T., Hall, G.E.M., and Cook, S.J. (Eds.), *Drift Exploration in Glaciated Terrain*, Geological Society, Special Publication 195, p. 45-68.

Maynard, D.E., 1989. Terrain analysis in exploration geochemistry. *Explore*, 65, 14-16.

McEneaney, M.B., Plouffe, A., McMartin, I., Campbell, J.E., Spittio, V.A., Paulen, R.C., Garrett, R.G., Hall, G.E.M., 2013. Till sampling and geochemical analytical protocols used by the Geological Survey of Canada. *Geochemistry, Exploration, Environment, Analysis*, 13, 285-301.

Miller, J.K., 1984. Model for elastic indicator trains in till. In: *Prospecting in areas of glaciated terrain*, A. Institute of Mining and Metallurgy, London, pp. 69-77.

Plouffe, A., Ferbey, T., Hashmi, S., and Ward, B.C., 2016. Till geochemistry and mineralogy: vectoring towards Cu porphyry deposits in British Columbia. *Canada Geochemistry: Exploration, Environment, Analysis*, <http://doi.org/10.1144/geochem2015-398>.

Proudfoot, D.N., and Allison, R.F., 1993a. Surficial geology of the Cusko River area (NTS 93C/9). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-18, 1:50,000 scale.

Proudfoot, D.N., and Allison, R.F., 1993b. Surficial geology of the Toll Mountain area (NTS 93C/16). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-19, 1:50,000 scale.

Proudfoot, D.N., Bobrowsky, P.T., and Meldrum, O.G., 1995. Drift exploration potential maps derived from terrain geology maps. In: Bobrowsky, P.T., Stribick, S.J., Newell, J.M., and Malysek, P.F. (Eds.), *Drift Exploration in the Canadian Cordillera*, British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Paper 1995-2, p. 23-31.

Saarnisto, M., 1990. An outline of glacial indicator tracing. In: Kujansuu, R., and Saarnisto, M. (Eds.), *Glacial Indicator Tracing*, Balkema, Rotterdam, 1-13.

Sacco, D.A., and Jackaman, W., 2015. Targeted geochemical and mineralogical surveys in the TREK project area, central British Columbia (parts of NTS 093B, C, F, G): year two. In: Geoscience BC Summary of Activities 2014, Geoscience BC, Report 2015-1, pp. 1-12.

Sacco, D., Ferbey, T., and Jackaman, W., 2014a. Basal till potential of the Cusko River map area (NTS 093C/9). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-06. Geoscience BC Map 2014-06-01, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014b. Basal till potential of the Cusko River map area (NTS 093B/12). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-07. Geoscience BC Map 2014-06-02, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014c. Basal till potential of the Marmot Lake map area (NTS 093B/13). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-06. Geoscience BC Map 2014-06-03, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014d. Basal till potential of the Toll Mountain map area (NTS 093C/16). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-09. Geoscience BC Map 2014-06-04, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014e. Basal till potential of the Suscha Creek map area (NTS 093F/1). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-10. Geoscience BC Map 2014-06-05, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014f. Basal till potential of the Cogitoko River map area (NTS 093G/04). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-11. Geoscience BC Map 2014-06-06, 1:50,000 scale.

Geological Survey Open File 2014-11; Geoscience BC Map 2014-06-06, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014g. Basal till potential of the Pelican Lake map area (NTS 093G/05). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-12. Geoscience BC Map 2014-06-07, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014h. Basal till potential of the Eucinka River map area (NTS 093F/08). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-13. Geoscience BC Map 2014-06-08, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014i. Basal till potential of the Chilako River map area (NTS 093G/12). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-14. Geoscience BC Map 2014-06-09, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014j. Basal till potential of the Hullat map area (NTS 093G/13). British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-15. Geoscience BC Map 2014-06-10, 1:50,000 scale.

Sacco, D.A., Jackaman, W., and Ferbey, T., 2014k. Targeted geochemical and mineralogical surveys in the TREK Project area, central British Columbia (parts of NTS 093B, C, F, G). In: Geoscience BC Summary of Activities 2013, Geoscience BC, Report 2014-1, pp. 19-34.


Shilts, W., 1993. Geological Survey of Canada's contributions to understanding the composition of glacial sediments. *Canadian Journal of Earth Sciences*, 30, p. 333-353.

Simard, G.J., Hora, Z.D., and Lefebvre, D.V. (Eds.), 1999. Selected British Columbia mineral deposit profiles, volume 3 - industrial minerals and gemstones. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1999-10, digital.

Spirito, W., McEneaney, M.B., Plouffe, A., McMartin, I., Campbell, J.E., Paulen, R.C., Garrett, R.G., and Hall, G.E.M., 2011. Till sampling and analytical protocols for GEM projects: from field to archive. Geological Survey of Canada, Open File 6850, 83 p.

Tipper, H.W., 1969. Geology, Anahim Lake, British Columbia. Geological Survey of Canada, Map 1220A, 1:252,440 scale.

Tipper, H.W., 1971. Surficial geology, Anahim Lake. Geological Survey of Canada, Map 1289A, 1:252,000 scale.



Recommended Citation:  
Sacco, D., Arnold, H., Ferbey, T., and Jackaman, W., 2017. Basal till potential of the Kushya River map area (NTS 093C/15), British Columbia. British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 2017-07, Geoscience BC Map 2017-02-06, 1:50,000 scale.